

Drug Facts

Active ingredient (in each tablet) Purpose

Aspirin 325 mg (NSAID*)..... Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains associated with

- | | | |
|---|---|---|
| <input type="checkbox"/> headache | <input type="checkbox"/> muscular aches | <input type="checkbox"/> minor arthritis pain |
| <input type="checkbox"/> backache | <input type="checkbox"/> common cold | <input type="checkbox"/> toothache |
| <input type="checkbox"/> menstrual cramps | | |

Temporarily reduces fever.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- | | | | |
|--|---|--|-------------------------------|
| <input type="checkbox"/> hives | <input type="checkbox"/> skin reddening | <input type="checkbox"/> facial swelling | <input type="checkbox"/> rash |
| <input type="checkbox"/> asthma (wheezing) | <input type="checkbox"/> blisters | <input type="checkbox"/> shock | |

If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- ☐ are age 60 or older
- ☐ have had stomach ulcers or bleeding problems
- ☐ take a blood thinning (anticoagulant) or steroid drug
- ☐ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- ☐ have 3 or more alcohol drinks every day while using this product
- ☐ take more or for a longer time than directed

Do not use

- ☐ if you have ever had an allergic reaction to any other pain reliever/fever reducer
- ☐ right before or after heart surgery
- ☐ if you are taking prescription drugs for gout, diabetes or arthritis

Ask a doctor before use if

- ☐ stomach bleeding warning applies to you
- ☐ you have a history of stomach problems such as heartburn
- ☐ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- ☐ you are taking a diuretic
- ☐ you have asthma

(Continued on opposite panel)

10202

**Drug Facts (continued)****Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ►

Drug Facts (continued)**Directions**

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- drink a full glass of water with each dose

Adults and children: (12 years and older)	Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.
--	---

Children under 12 years:	Do not give to children under 12 years of age.
---------------------------------	--

Other information

- read all product information before using
- store at room temperature 15°-30°C (59°-86°F)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Questions or comments? 1-800-634-7680